

Transmembrane Protein 43 Rabbit mAb

Catalog # AP76183

Product Information

ApplicationWB, IHC-PPrimary AccessionQ9BTV4ReactivityHuman, RatHostRabbit

Clonality Monoclonal Antibody

Calculated MW 44876

Additional Information

Gene ID 79188

Other Names TMEM43

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name TMEM43

Function May have an important role in maintaining nuclear envelope structure by

organizing protein complexes at the inner nuclear membrane. Required for retaining emerin at the inner nuclear membrane (By similarity). Plays a role in the modulation of innate immune signaling through the cGAS-STING pathway by interacting with RNF26 (PubMed:32614325). In addition, functions as a critical signaling component in mediating NF-kappa-B activation by acting downstream of EGFR and upstream of CARD10 (PubMed:27991920). Contributes to passive conductance current in cochlear glia-like supporting cells, mediated by gap junctions and necessary for hearing and speech

discrimination (PubMed:34050020).


Cellular Location Endoplasmic reticulum membrane. Nucleus inner membrane; Multi-pass


membrane protein. Cell membrane Note=Retained in the inner nuclear membrane through interaction with EMD and A- and B-lamins. The N- and C-termini are oriented towards the nucleoplasm. The majority of the hydrophilic domain resides in the endoplasmic reticulum lumen (By

similarity).

Highest expression in placenta. Also found at lower levels in heart, ovary, spleen, small intestine, thymus, prostate and testis.

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.