

TSG101 Rabbit mAb

Catalog # AP76209

Product Information

Application	WB, IP
Primary Accession	Q99816
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	43944


Additional Information

Gene ID	7251
Other Names	TSG101
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	Liquid

Protein Information

Name	TSG101
Function	Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55. May be involved in cell growth and differentiation. Acts as a negative growth regulator. Involved in the budding of many viruses through an interaction with viral proteins that contain a late-budding motif P-[ST]-A-P. This interaction is essential for viral particle budding of numerous retroviruses. Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed: 22660413). It may also play a role in the extracellular release of microvesicles that differ from the exosomes (PubMed: 22315426).
Cellular Location	Cytoplasm. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Late endosome membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Midbody, Midbody ring. Nucleus. Note=Mainly cytoplasmic. Membrane-associated when active and soluble when inactive. Nuclear localization is cell cycle-dependent. Interaction with CEP55 is required for localization to the midbody during cytokinesis
Tissue Location	Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.