

Vinculin Rabbit mAb

Catalog # AP76249

Product Information

Application	WB, IP, ICC
Primary Accession	P18206
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	123799

Additional Information

Gene ID	7414
Other Names	VCL
Dilution	WB~~1/500-1/1000 IP~~N/A ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information



Name	VCL
Function	Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250 UniProtKB:P12003}; Cytoplasmic side {ECO:0000250 UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250 UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250 UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250 UniProtKB:Q64727}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250 UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250 UniProtKB:Q64727}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions

{ECO:0000250|UniProtKB:P12003}

Tissue Location

Metavinculin is muscle-specific.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.