

Vinculin Rabbit mAb

Catalog # AP76250

Product Information

Application	WB, IHC-P
Primary Accession	P18206
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	123799

Additional Information

Gene ID	7414
Other Names	VCL
Dilution	WB~1/500-1/1000 IHC-P~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.



Protein Information

Name	VCL
Function	Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250 UniProtKB:P12003}; Cytoplasmic side {ECO:0000250 UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250 UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250 UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250 UniProtKB:Q64727}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250 UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250 UniProtKB:Q64727}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions

Tissue Location

Metavinculin is muscle-specific.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.