10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

VPS41 Rabbit mAb

Catalog # AP76254

Product Information

Application	WB
Primary Accession	P49754
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	98566

Additional Information

Gene ID	27072
Other Names	VPS41
Dilution	WB~~1:500-1:1000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	VPS41
Function	Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Acts as a component of the HOPS endosomal tethering complex. This complex is proposed to be involved in the Rab5- to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes (PubMed:23351085, PubMed:33851776). Involved in homotypic vesicle fusions between late endosomes and in heterotypic fusions between late endosomes and lysosomes implicated in degradation of endocytosed cargo (PubMed:23167963, PubMed:25445562, PubMed:25908847, PubMed:9159129). Required for fusion of autophagosomes with lysosomes (PubMed:25783203, PubMed:37821429). Links the HOPS complex to

	endosomal Rab7 via its association with RILP and to lysosomal membranes via its association with ARL8B, suggesting that these interactions may bring the compartments to close proximity for fusion (PubMed:21802320, PubMed:25445562, PubMed:25908847). Involved in the direct trans-Golgi network to late endosomes transport of lysosomal membrane proteins independently of HOPS (PubMed:23322049). Involved in sorting to the regulated secretory pathway presumably implicating the AP-3 adapter complex (By similarity). May play a role in HOPS-independent function in the regulated secretory pathway (PubMed:24210660).
Cellular Location	Endosome membrane; Peripheral membrane protein. Late endosome membrane; Peripheral membrane protein. Early endosome membrane; Peripheral membrane protein. Lysosome membrane; Peripheral membrane protein. Golgi apparatus, trans- Golgi network. Cytoplasmic vesicle, clathrin-coated vesicle. Cytoplasm, cytosol
Tissue Location	Expressed in cerebral cortex and cerebellum. Highly expressed in Purkinje cells.

Background


Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act in part as a core component of the putative HOPS endosomal tethering complex is proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes (PubMed:23351085). Involved in homotypic vesicle fusions between late endosomes and in heterotypic fusions between late endosomes and lysosomes implicated in degradation of endocytosed cargo (PubMed:9159129, PubMed:23167963, PubMed:25445562, PubMed:25908847). Required for fusion of autophagosomes with lysosomes (PubMed:25783203). May link the HOPS complex to endosomal Rab7 via its association with RILP and to lysosomal membranes via its association with ARL8B, suggesting that these interactions may bring the compartments to close proximity for fusion (PubMed:25445562, PubMed:25908847). Involved in the direct trans-Golgi network to late endosomes transport of lysosomal membrane proteins independently of HOPS (PubMed:23322049). Involved in sorting to the regulated secretory pathway presumably implicating the AP-3 adaptor complex . May play a role in HOPS-independent function in the regulated secretory pathway (PubMed:24210660).

Images



Western blot analysis of VPS41 in rat heart lysates using VPS41 antibody.

Western blot analysis of VPS41 in Hela, 293T, mouse brain lysates using VPS41 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.