10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Phospho-YAP1 (Ser127) Rabbit mAb

Catalog # AP76266

Product Information

Application WB, IHC-P Primary Accession P46937

Reactivity Human, Mouse, Rat

Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 54462

Additional Information

Gene ID 10413

Other Names YAP1

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format Liquid

Protein Information

Name YAP1 (HGNC:16262)

Synonyms YAP65

Function Transcriptional regulator with dual roles as a coactivator and corepressor.

Critical downstream regulatory target in the Hippo signaling pathway, crucial for organ size control and tumor suppression by restricting proliferation and

promoting apoptosis (PubMed: 17974916, PubMed: 18280240,

PubMed: 18579750, PubMed: 21364637, PubMed: 30447097). The Hippo signaling pathway core involves a kinase cascade featuring STK3/MST2 and STK4/MST1, along with its regulatory partner SAV1, which phosphorylates and activates LATS1/2 in complex with their regulatory protein, MOB1. This activation leads to the phosphorylation and inactivation of the YAP1

oncoprotein and WWTR1/TAZ (PubMed:<u>18158288</u>). Phosphorylation of YAP1 by LATS1/2 prevents its nuclear translocation, thereby regulating the expression of its target genes (PubMed:<u>18158288</u>, PubMed:<u>26598551</u>,

PubMed:34404733). The transcriptional regulation of gene expression requires TEAD transcription factors and modulates cell growth,

anchorage-independent growth, and induction of epithelial- mesenchymal transition (EMT) (PubMed: 18579750). Plays a key role in tissue tension and 3D tissue shape by regulating the cortical actomyosin network, acting via

ARHGAP18, a Rho GTPase activating protein that suppresses F-actin polymerization (PubMed: 25778702). It also suppresses ciliogenesis by acting as a transcriptional corepressor of TEAD4 target genes AURKA and PLK1

(PubMed: <u>25849865</u>). In conjunction with WWTR1, regulates TGFB1-dependent SMAD2 and SMAD3 nuclear accumulation (By similarity). Synergizes with WBP2 to enhance PGR activity (PubMed: <u>16772533</u>).


Cellular Location

Cytoplasm. Nucleus. Cell junction, tight junction {ECO:0000250|UniProtKB:A0A8C0NGY6}. Cell membrane. Note=Both phosphorylation and cell density can regulate its subcellular localization (PubMed:18158288, PubMed:20048001). Phosphorylation sequesters it in the cytoplasm by inhibiting its translocation into the nucleus (PubMed:18158288, PubMed:20048001, PubMed:34404733). At low density, predominantly nuclear and is translocated to the cytoplasm at high density (PubMed:18158288, PubMed:20048001, PubMed:25849865). PTPN14 induces translocation from the nucleus to the cytoplasm (PubMed:22525271). In the nucleus, phosphorylation by PRP4K induces nuclear exclusion (PubMed:29695716). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity) Localizes to the cytoplasm and tight junctions following interaction with AMOT isoform 1 (PubMed:21205866). Localizes to tight junctions following interaction with AMOTL2 (By similarity). Translocates to the nucleus in the presence of SNAIL1 (By similarity). Found at the cell membrane in keratinocytes in response to mechanical strain (PubMed:31835537). {ECO:0000250|UniProtKB:A0A8C0NGY6, ECO:0000250|UniProtKB:P46938. ECO:0000269 | PubMed:18158288, ECO:0000269 | PubMed:20048001, ECO:0000269 | PubMed:21205866, ECO:0000269 | PubMed:22525271, ECO:0000269 | PubMed:25849865, ECO:0000269 | PubMed:29695716, ECO:0000269 | PubMed:31835537, ECO:0000269 | PubMed:34404733 }

Tissue Location

Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.