10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TIM 1 Rabbit mAb

Catalog # AP76285

Product Information

Application	WB
Primary Accession	Q96D42
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	39250

Additional Information

Gene ID	26762
Other Names	HAVCR1
Dilution	WB~~1:2000-1:5000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	HAVCR1
Synonyms	KIM1, TIM1, TIMD1
Function	Phosphatidylserine receptor that plays an important functional role in regulatory B-cell homeostasis including generation, expansion and suppressor functions (PubMed:18082433, PubMed:18414680). Involved in efferocytosis, the process by which apoptotic cells are removed, by binding to phosphatidylserine present on apoptotic cells and promoting their engulfment (PubMed:18082433). Confers phagocytic ability on injured kidney epithelial cells, allowing them to bind to and internalize apoptotic bodies and necrotic debris following kidney injury (PubMed:17471468, PubMed:18414680). The ectodomain binds to the surface of apoptotic kidney epithelial cells via phosphatidylserine and oxidized phospholipids exposed on the cell membrane of the apoptotic cells (PubMed:18414680). HAVCR1-expressing cells are also capable of phagocytosing Gram-negative and Gram-positive bacteria (PubMed:18414680). As P-selectin/SELPLG ligand,

	plays a specialized role in activated but not naive T-cell trafficking during inflammatory responses (PubMed:24703780). Controls thereby T-cell accumulation in the inflamed central nervous system (CNS) and the induction of autoimmune disease (PubMed:24703780). Acts as a regulator of T-cell proliferation (By similarity). Also regulates expression of various anti-inflammatory cytokines and co-inhibitory ligands including IL10 (By similarity).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Cell projection, phagocytic cup. Cytoplasmic vesicle, phagosome
Tissue Location	Widely expressed, with highest levels in kidney and testis. Expressed by activated CD4+ T-cells during the development of helper T-cells responses.


Background

May play a role in T-helper cell development and the regulation of asthma and allergic diseases. Receptor for TIMD4 (By similarity). In case of human hepatitis A virus (HHAV) infection, functions as a cell-surface receptor for the virus. May play a role in kidney injury and repair.

Images



Western blot analysis of TIM 1 in Raw264.7, PC-12 lysates using Hepatitis A Virus ular Receptor 1 antibody.



Western blot analysis of TIM 1 in A549, MCF-7, U251, HL-60 lysates using TIM 1 antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.