

# Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb

Catalog # AP76334

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB                     |
| Primary Accession | <a href="#">P13639</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 95338                  |

## Additional Information

---

|             |  |
|-------------|--|
| Gene ID     | 1938   |
| Other Names | EEF2   |
| Dilution    | WB~~1/500-1/1000   |
| Format      | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.    |
| Storage     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |


## Protein Information

---

|                   |  |
|-------------------|--|
| Name              | EEF2   |
| Synonyms          | EF2  |
| Function          | Catalyzes the GTP-dependent ribosomal translocation step during translation elongation (PubMed: <a href="#">26593721</a> ). During this step, the ribosome changes from the pre-translocational (PRE) to the post- translocational (POST) state as the newly formed A-site-bound peptidyl- tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively (PubMed: <a href="#">26593721</a> ). Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome (PubMed: <a href="#">26593721</a> ). |
| Cellular Location | Cytoplasm. Nucleus. Note=Phosphorylation by CSK promotes cleavage and SUMOylation-dependent nuclear translocation of the C- terminal cleavage product.   |

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.