

Phospho-Hormone Sensitive Lipase (Ser855) Rabbit mAb

Catalog # AP76344

Product Information

Application	WB
Primary Accession	Q05469
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	116598

Additional Information

Gene ID	3991
Other Names	LIPE
Dilution	WB~~1/500-1/1000
Format	Liquid


Protein Information

Name	LIPE
Function	<p>Lipase with broad substrate specificity, catalyzing the hydrolysis of triacylglycerols (TAGs), diacylglycerols (DAGs), monoacylglycerols (MAGs), cholesteryl esters and retinyl esters (PubMed:15716583, PubMed:15955102, PubMed:19800417, PubMed:8812477). Shows a preferential hydrolysis of DAGs over TAGs and MAGs and preferentially hydrolyzes the fatty acid (FA) esters at the sn-3 position of the glycerol backbone in DAGs (PubMed:19800417). Preferentially hydrolyzes FA esters at the sn-1 and sn-2 positions of the glycerol backbone in TAGs (By similarity). Catalyzes the hydrolysis of 2-arachidonoylglycerol, an endocannabinoid and of 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor (By similarity). In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free cholesterol for steroid hormone production (By similarity).</p>
Cellular Location	<p>Cell membrane. Membrane, caveola. Cytoplasm, cytosol. Lipid droplet {ECO:0000250 UniProtKB:P54310}. Note=Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation (PubMed:17026959). Phosphorylation by AMPK reduces its translocation towards the lipid droplets (By similarity) {ECO:0000250 UniProtKB:P54310, ECO:0000269 PubMed:17026959}</p>

Tissue Location

Testis..

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.