

14-3-3 gamma Rabbit mAb

Catalog # AP76372

Product Information

Application	WB
Primary Accession	P61981
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	28303

Additional Information

Gene ID	7532
Other Names	YWHAG
Dilution	WB~~1/500-1/1000
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	YWHAG (HGNC:12852)
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binding generally results in the modulation of the activity of the binding partner (PubMed: 16511572). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed: 36732624). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed: 29121962). Reduces keratinocyte intercellular adhesion, via interacting with PKP1 and sequestering it in the cytoplasm, thereby reducing its incorporation into desmosomes (PubMed: 29678907). Plays a role in mitochondrial protein catabolic process (also named MALM) that promotes the degradation of damaged proteins inside mitochondria (PubMed: 22532927).
Cellular Location	Cytoplasm, cytosol. Mitochondrion matrix. Note=Translocates to the

mitochondrial matrix following induction of MALM (mitochondrial protein catabolic process).

Tissue Location

Highly expressed in brain, skeletal muscle, and heart.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.