

14-3-3 gamma Rabbit mAb

Catalog # AP76372

Product Information

Application	WB
Primary Accession	<u>P61981</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	28303

Additional Information

Gene ID	7532
Other Names	YWHAG
Dilution	WB~~1/500-1/1000
Format	Liquid

Protein Information

Name	YWHAG (<u>HGNC:12852</u>)
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: <u>15696159</u> , PubMed: <u>16511572</u> , PubMed: <u>36732624</u>). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: <u>15696159</u> , PubMed: <u>16511572</u> , PubMed: <u>36732624</u>). Binding generally results in the modulation of the activity of the binding partner (PubMed: <u>16511572</u>). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed: <u>36732624</u>). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed: <u>29121962</u>). Reduces keratinocyte intercellular adhesion, via interacting with PKP1 and sequestering it in the cytoplasm, thereby reducing its incorporation into desmosomes (PubMed: <u>29678907</u>). Plays a role in mitochondrial protein catabolic process (also named MALM) that promotes the degradation of damaged proteins inside mitochondria (PubMed: <u>22532927</u>).
Cellular Location	Cytoplasm, cytosol. Mitochondrion matrix. Note=Translocates to the mitochondrial matrix following induction of MALM (mitochondrial protein catabolic process).
Tissue Location	Highly expressed in brain, skeletal muscle, and heart.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.