

5T4 Rabbit mAb

Catalog # AP76373

Product Information

Application	WB, IP
Primary Accession	Q13641
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	46032

Additional Information

Gene ID	7162
Other Names	TPBG
Dilution	WB~~1:1000-1:5000 IP~~1:50-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	TPBG
Synonyms	5T4
Function	May function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization.
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed by all types of trophoblasts as early as 9 weeks of development. Specific for trophoblastic cells except for amniotic epithelium. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma


Background

5T4 is a 72 kDa transmembrane glycoprotein that exhibits restricted expression in human and mouse adult tissues, is upregulated on many carcinomas and tumour expression correlates with poorer clinical outcome in ovarian, gastric and colorectal cancers. mES cells lack cell surface expression of the 5T4 antigen, and both protein and mRNA are rapidly upregulated following induction of differentiation. This proteins expression correlates with downregulation of OCT-4 and Tra-1-60.

Images



Western blot analysis of 5T4 in mouse brain, rat brain lysates using 5T4 antibody.



Western blot analysis of 5T4 in 293T, C2C12 lysates using 5T4 antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.