

ADK Rabbit mAb

Catalog # AP76381

Product Information

Application	WB, IHC-P, FC
Primary Accession	P55263
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	40545

Additional Information

Gene ID	132
Other Names	ADK
Dilution	WB~~1:500-1:1000 IHC-P~~N/A FC~~1:200-1:500
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ADK {ECO:0000303 PubMed:19635462, ECO:0000312 HGNC:HGNC:257}
Function	Adenosine kinase that mediates the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner: catalyzes phosphorylation of both unmodified and modified adenosines (PubMed: 21963049 , PubMed: 40840445 , PubMed: 6246102 , PubMed: 8577746 , PubMed: 9070863). Plays a key role in the detoxification of modified adenosines containing N(6)-methylated adenine (m6A) post- transcriptional modification (PubMed: 40840445). Modified nucleosides are derived from the degradation of RNAs (mRNAs, rRNAs and tRNAs) and possess intrinsic cytotoxicity and must be cleared to prevent metabolic dysfunction (PubMed: 40840445). Catalyzes the phosphorylation of the free cytosolic methylated adenosine nucleotides N(6)-methyladenosine (m6A), N(6),N(6)-dimethyladenosine (m6,6A) and N(6)- isopentenyladenosine (i6A) into adenosine monophosphate (AMP) intermediates that are further detoxified by MAPDA/ADAL (PubMed: 40840445).

Cellular Location


Cytoplasm, cytosol. [Isoform 2]: Cytoplasm

Tissue Location


Widely expressed. Highest level in placenta, liver, muscle and kidney.

Background


This gene encodes an enzyme which catalyzes the transfer of the gamma-phosphate from ATP to adenosine, thereby serving as a regulator of concentrations of both extracellular adenosine and intracellular adenine nucleotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune systems and inhibitors of the enzyme could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as anti-inflammatory agents. Multiple transcript variants encoding different isoforms have been found for this gene.

Images

Western blot analysis of ADK in PC-12, Raw264.7 lysates using ADK antibody.



Western blot analysis of ADK in HeLa, A549, HL-60, U251, U87-MG lysates using ADK antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using ADK antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.