

AMPK gamma 1 Rabbit mAb

Catalog # AP76387

Product Information

Application WB, IP, ICC
Primary Accession P54619
Reactivity Human
Rabbit

Clonality Monoclonal Antibody

Calculated MW 37579

Additional Information

Gene ID 5571

Other Names PRKAG1

Dilution WB~~1/500-1/1000 IP~~N/A ICC~~N/A

Format Liquid

Protein Information

Name PRKAG1

Function AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy

sensor protein kinase that plays a key role in regulating cellular energy metabolism (PubMed:21680840, PubMed:24563466). In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation

(PubMed:<u>21680840</u>, PubMed:<u>24563466</u>). AMPK acts via direct

phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators (PubMed: <u>21680840</u>,

PubMed: 24563466). Also acts as a regulator of cellular polarity by remodeling

the actin cytoskeleton; probably by indirectly activating myosin


(PubMed:<u>21680840</u>, PubMed:<u>24563466</u>). Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1


or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits (PubMed: <u>21680840</u>,

PubMed: 24563466). ADP also stimulates phosphorylation, without stimulating

already phosphorylated catalytic subunit (PubMed: 21680840,

PubMed: <u>24563466</u>). ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive (PubMed: <u>21680840</u>, PubMed: <u>24563466</u>).





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.