

ABCG2 Rabbit mAb

Catalog # AP76407

Product Information

Application	WB
Primary Accession	Q9UNQ0
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	72314

Additional Information

Gene ID	9429
Other Names	ABCG2
Dilution	WB~~1:1000-1:5000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ABCG2 (HGNC:74)
Synonyms	ABCP, BCRP, BCRP1, MXR
Function	Broad substrate specificity ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes a wide variety of physiological compounds, dietary toxins and xenobiotics from cells (PubMed: 11306452 , PubMed: 12958161 , PubMed: 19506252 , PubMed: 20705604 , PubMed: 28554189 , PubMed: 30405239 , PubMed: 31003562). Involved in porphyrin homeostasis, mediating the export of protoporphyrin IX (PPIX) from both mitochondria to cytosol and cytosol to extracellular space, it also functions in the cellular export of heme (PubMed: 20705604 , PubMed: 23189181). Also mediates the efflux of sphingosine-1-P from cells (PubMed: 20110355). Acts as a urate exporter functioning in both renal and extrarenal urate excretion (PubMed: 19506252 , PubMed: 20368174 , PubMed: 22132962 , PubMed: 31003562 , PubMed: 36749388). In kidney, it also functions as a physiological exporter of the uremic toxin indoxyl sulfate (By

similarity). Also involved in the excretion of steroids like estrone 3-sulfate/E1S, 3beta-sulfooxy-androst-5-en-17-one/DHEAS, and other sulfate conjugates (PubMed:[12682043](#), PubMed:[28554189](#), PubMed:[30405239](#)). Mediates the secretion of the vitamins riboflavin and biotin into milk (By similarity). Involved in the excretion of the riboflavin-derived compound lumichrome into the intestinal lumen and in its secretion into milk (PubMed:[39337371](#)). Extrudes pheophorbide a, a phototoxic porphyrin catabolite of chlorophyll, reducing its bioavailability (By similarity). Plays an important role in the exclusion of xenobiotics from the brain (Probable). It confers to cells a resistance to multiple drugs and other xenobiotics including mitoxantrone, pheophorbide, camptothecin, methotrexate, azidothymidine, and the anthracyclines daunorubicin and doxorubicin, through the control of their efflux (PubMed:[11306452](#), PubMed:[12477054](#), PubMed:[15670731](#), PubMed:[18056989](#), PubMed:[31254042](#)). In placenta, it limits the penetration of drugs from the maternal plasma into the fetus (By similarity). May play a role in early stem cell self-renewal by blocking differentiation (By similarity). In inflammatory macrophages, exports itaconate from the cytosol to the extracellular compartment and limits the activation of TFEB-dependent lysosome biogenesis involved in antibacterial innate immune response.

Cellular Location

Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Mitochondrion membrane; Multi-pass membrane protein. Note=Enriched in membrane lipid rafts


Tissue Location

Highly expressed in placenta (PubMed:[9850061](#)). Low expression in small intestine, liver and colon (PubMed:[9861027](#)) Expressed in brain (at protein level) (PubMed:[12958161](#))

Background

The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene.

Images



Western blot analysis of BCRP/ABCG2 in HepG2, U251, HL-60 lysates using ABCG2 antibody.