

IGFBP3 Antibody (S183)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7641D

Product Information

Application	WB, E
Primary Accession	P17936
Other Accession	P16611 , P20959
Reactivity	Human, Hamster, Rat, Mouse
Predicted	Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31674
Antigen Region	162-189

Additional Information

Gene ID	3486
Other Names	Insulin-like growth factor-binding protein 3, IBP-3, IGF-binding protein 3, IGFBP-3, IGFBP3, IBP3
Target/Specificity	This IGFBP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 162-189 amino acids from human IGFBP3.
Dilution	WB~~1 : 1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	IGFBP3 Antibody (S183) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IGFBP3
Synonyms	IBP3
Function	Multifunctional protein that plays a critical role in regulating the availability

of IGFs such as IGF1 and IGF2 to their receptors and thereby regulates IGF-mediated cellular processes including proliferation, differentiation, and apoptosis in a cell-type specific manner (PubMed:[10874028](#), PubMed:[19556345](#)). Also exhibits IGF- independent antiproliferative and apoptotic effects mediated by its receptor TMEM219/IGFBP-3R (PubMed:[20353938](#)). Inhibits the positive effect of humanin on insulin sensitivity (PubMed:[19623253](#)). Promotes testicular germ cell apoptosis (PubMed:[19952275](#)). Acts via LRP- 1/ α 2M receptor, also known as TGF-beta type V receptor, to mediate cell growth inhibition independent of IGF1 (PubMed:[9252371](#)). Mechanistically, induces serine-specific dephosphorylation of IRS1 or IRS2 upon ligation to its receptor, leading to the inhibitory cascade (PubMed:[15371331](#)). In the nucleus, interacts with transcription factors such as retinoid X receptor- α /RXRA to regulate transcriptional signaling and apoptosis (PubMed:[10874028](#)).

Cellular Location

Secreted. Nucleus

Tissue Location

Expressed by most tissues. Present in plasma.


Background

IGFBP3 is a member of the insulin-like growth factor binding protein (IGFBP) family with an IGFBP domain and a thyroglobulin type-I domain. This protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors.

References

Muzumdar,R.H., Diabetes 55 (10), 2788-2796 (2006) Novosyadlyy,R., Growth Horm. IGF Res. 15 (5), 313-323 (2005)

Images



Anti-IGFBP-3-S183 Antibody at 1:1000 dilution + human liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.