

# EBP1 Rabbit mAb

Catalog # AP76474

### **Product Information**

ApplicationWB, IHC-P, IPPrimary AccessionQ9UQ80

**Reactivity** Human, Mouse, Rat, Hamster

**Host** Rabbit

**Clonality** Monoclonal Antibody

Calculated MW 43787

## **Additional Information**

**Gene ID** 5036

Other Names PA2G4

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A IP~~1/20

Format Liquid

#### **Protein Information**

Name PA2G4

Synonyms EBP1

**Function** May play a role in a ERBB3-regulated signal transduction pathway. Seems be

involved in growth regulation. Acts a corepressor of the androgen receptor (AR) and is regulated by the ERBB3 ligand neuregulin-1/heregulin (HRG). Inhibits transcription of some E2F1- regulated promoters, probably by recruiting histone acetylase (HAT) activity. Binds RNA. Associates with 28S, 18S and 5.8S mature rRNAs, several rRNA precursors and probably U3 small nucleolar RNA. May be involved in regulation of intermediate and late steps of rRNA processing. May be involved in ribosome assembly. Mediates capindependent translation of specific viral IRESs (internal ribosomal entry site) (By similarity). Regulates cell proliferation, differentiation, and survival. Isoform 1 suppresses apoptosis whereas isoform 2 promotes cell

differentiation (By similarity).

**Cellular Location** [Isoform 1]: Cytoplasm. Nucleus, nucleolus Note=Translocates to the nucleus


upon treatment with HRG Phosphorylation at Ser-361 by PKC/PRKCD


regulates its nucleolar localization.


**Tissue Location** Isoform 2 is undetectable whereas isoform 1 is strongly expressed in cancer

cells (at protein level). Isoform 1 and isoform 2 are widely expressed, including heart, brain, lung, pancreas, skeletal muscle, kidney, placenta and

# **Images**







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.