

Elongation factor 2 Rabbit mAb

Catalog # AP76477

Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	P13639
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	95338





Additional Information

Gene ID	1938
Other Names	EEF2
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A
Format	Liquid

Protein Information

Name	EEF2
Synonyms	EF2
Function	Catalyzes the GTP-dependent ribosomal translocation step during translation elongation (PubMed: 26593721). During this step, the ribosome changes from the pre-translocational (PRE) to the post- translocational (POST) state as the newly formed A-site-bound peptidyl- tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively (PubMed: 26593721). Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome (PubMed: 26593721).
Cellular Location	Cytoplasm. Nucleus. Note=Phosphorylation by CSK promotes cleavage and SUMOylation-dependent nuclear translocation of the C- terminal cleavage product.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.