

Lactoferrin (LTF) Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7647c

Product Information

Application IHC-P, WB, E **Primary Accession** P02788

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalIsotypeRabbit IgGCalculated MW78182Antigen Region219-248

Additional Information

Gene ID 4057

Other Names Lactotransferrin, Lactoferrin, 3421-, Growth-inhibiting protein 12,

Talalactoferrin, Lactoferricin-H, Lfcin-H, Kaliocin-1, Lactoferroxin-A,

Lactoferroxin-B, Lactoferroxin-C, LTF, GIG12, LF

Target/SpecificityThis Lactoferrin (LTF) antibody is generated from rabbits immunized with a

KLH conjugated synthetic peptide between 219-248 amino acids from the

Central region of human Lactoferrin (LTF).

Dilution IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Lactoferrin (LTF) Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name LTF (HGNC:6720)

Synonyms GIG12, LF

Function Transferrins are iron binding transport proteins which can bind two Fe(3+)

ions in association with the binding of an anion, usually bicarbonate.

Cellular Location

[Isoform 1]: Secreted. Cytoplasmic granule. Note=Secreted into most exocrine fluids by various endothelial cells Stored in the secondary granules of neutrophils


Tissue Location

High levels are found in saliva and tears, intermediate levels in serum and plasma, and low levels in urine. In kidney, detected in the distal collecting tubules in the medulla but not in the cortical region or in blood vessels. Detected in peripheral blood neutrophils (at protein level). Isoform 1 and isoform DeltaLf are expressed in breast, prostate, spleen, pancreas, kidney, small intestine, lung, skeletal muscle, uterus, thymus and fetal liver Isoform 1 is expressed in brain, testis and peripheral blood leukocytes; isoform DeltaLf is barely detectable in these tissues Isoform DeltaLf is expressed in placenta, liver and ovary; isoform 1 is barely detectable in these tissues. In kidney, isoform 1 is expressed at high levels in the collecting tubules of the medulla but at very low levels in the cortex.


Background

This protein is a member of the transferrin family of metal-binding proteins found in milk and other secretory fluids and also in blood. It shows multifunctional properties of which the bacteriostatic and bactericidal effects are the best known.

Images




Formalin-fixed and paraffin-embedded prostata carcinoma tissue reacted with LTF Antibody (Center)(Cat.#AP7647c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Western blot analysis of anti-LTF Antibody (Center) Pab (Cat.#AP7647c) in mouse heart tissue lysates (35ug/lane). LTF(arrow) was detected using the purified Pab (1:60 dilution).

Western blot analysis of anti-LTF Antibody (Center) Pab (Cat.#AP7647c) in 293 cell line lysates transiently transfected with the LTF gene (2ug/lane). LTF(arrow) was detected using the purified Pab (1:60 dilution).



Citations

• Tumor microenvironment-derived proteins dominate the plasma proteome response during breast cancer induction and progression.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.