

EYA2 Rabbit mAb

Catalog # AP76490

Product Information

Application	WB, IP
Primary Accession	O00167
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	59232


Additional Information

Gene ID	2139
Other Names	EYA2
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	Liquid

Protein Information

Name	EYA2
Synonyms	EAB1
Function	Functions both as protein phosphatase and as transcriptional coactivator for SIX1, and probably also for SIX2, SIX4 and SIX5 (PubMed: 12500905 , PubMed: 23435380). Tyrosine phosphatase that dephosphorylates 'Tyr-142' of histone H2AX (H2AXY142ph) and promotes efficient DNA repair via the recruitment of DNA repair complexes containing MDC1. 'Tyr-142' phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress (PubMed: 19351884). Its function as histone phosphatase may contribute to its function in transcription regulation during organogenesis. Plays an important role in hypaxial muscle development together with SIX1 and DACH2; in this it is functionally redundant with EYA1 (PubMed: 12500905).
Cellular Location	Cytoplasm. Nucleus Note=Retained in the cytoplasm via interaction with GNAZ and GNAI2 (PubMed:10906137). Interaction with SIX1, SIX2, SIX4 or SIX5 is required for translocation to the nucleus (PubMed:10906137, PubMed:12500905).
Tissue Location	Highest expression in muscle with lower levels in kidney, placenta, pancreas, brain and heart

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.