

GRK2 Rabbit mAb

Catalog # AP76516

Product Information

Application WB, IP
Primary Accession P25098
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 79574

Additional Information

Gene ID 156

Other Names GRK2

Dilution WB~~1/500-1/1000 IP~~N/A

Format Liquid

Protein Information

Name GRK2 (<u>HGNC:289</u>)

Synonyms ADRBK1, BARK, BARK1

Function Specifically phosphorylates the agonist-occupied form of the

beta-adrenergic and closely related receptors, probably inducing a

desensitization of them (PubMed:<u>19715378</u>). Key regulator of LPAR1 signaling (PubMed:<u>19306925</u>). Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor (PubMed:<u>19306925</u>). Desensitizes

LPAR1 and LPAR2 in a phosphorylation- independent manner (PubMed: 19306925). Positively regulates ciliary smoothened

(SMO)-dependent Hedgehog (Hh) signaling pathway by facilitating the trafficking of SMO into the cilium and the stimulation of SMO activity (By similarity). Inhibits relaxation of airway smooth muscle in response to blue

light (PubMed: 30284927).


Cellular Location Cytoplasm {ECO:0000250 | UniProtKB:P26817}. Cell membrane

{ECO:0000250 | UniProtKB:P21146}. Postsynapse {ECO:0000250 | UniProtKB:P26817}. Presynapse

{ECO:0000250 | UniProtKB:P26817}

Tissue Location Expressed in peripheral blood leukocytes.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.