

hHR23A Rabbit mAb

Catalog # AP76525

Product Information

Primary Accession	P54725
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	39609


Additional Information

Gene ID	5886
Other Names	RAD23A
Format	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	RAD23A (HGNC:9812)
Function	Multifunctional protein that participates in histone H4K20 demethylation, DNA repair, ubiquitin-dependent protein degradation, transcriptional regulation, and viral replication (PubMed:12643283, PubMed:14621999, PubMed:15321727, PubMed:20614012, PubMed:32209475, PubMed:9372924). Specifically demethylates mono-, di- and trimethylated 'Lys-20' of histone H4 (H4K20me1, H4K20me2, H4K20me3, respectively) into unmethylated forms. Activates the transcription of coding genes by demethylating H4K20me1 and the transcription of repetitive elements by demethylating H4K20me3 (PubMed:32209475). Involved in modulation of proteasomal degradation. Binds to 'Lys-48'-linked polyubiquitin chains in a length-dependent manner and with a lower affinity to 'Lys-63'- linked polyubiquitin chains. Proposed to be capable to bind simultaneously to the 26S proteasome and to polyubiquitinated substrates and to deliver ubiquitinated proteins to the proteasome (PubMed:12643283, PubMed:14621999, PubMed:15321727). Involved in nucleotide excision repair, is considered functionally equivalent to RAD23B in global genome nucleotide excision repair (GG-NER) through its association with XPC. In vitro, the XPC-RAD23A complex demonstrates NER activity (PubMed:9372924). Can stabilize XPC (By similarity).
Cellular Location	Nucleus.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.