

Methionine Aminopeptidase 2 Rabbit mAb

Catalog # AP76589

Product Information

Primary Accession	P50579
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	52892



Additional Information

Gene ID	10988
Other Names	METAP2
Format	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	METAP2 {ECO:0000255 HAMAP-Rule:MF_03175}
Synonyms	MNPEP, P67EIF2
Function	Cotranslationally removes the N-terminal methionine from nascent proteins. The N-terminal methionine is often cleaved when the second residue in the primary sequence is small and uncharged (Met- Ala-, Cys, Gly, Pro, Ser, Thr, or Val). The catalytic activity of human METAP2 toward Met-Val peptides is consistently two orders of magnitude higher than that of METAP1, suggesting that it is responsible for processing proteins containing N-terminal Met-Val and Met-Thr sequences in vivo.
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03175, ECO:0000269 PubMed:21537465}. Note=About 30% of expressed METAP2 associates with polysomes

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.