

Moesin Rabbit mAb

Catalog # AP76596

Product Information

Application WB, IHC-P, IHC-F, IP, ICC

Primary Accession <u>P26038</u>

Reactivity Human, Mouse, Rat

Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 67820

Additional Information

Gene ID 4478

Other Names MSN

Dilution WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name MSN (HGNC:7373)

Function Ezrin-radixin-moesin (ERM) family protein that connects the actin

cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:10212266). Once phosphorylated on its C-terminal threonine, moesin is activated leading

to interaction with F-actin and cytoskeletal rearrangement

(PubMed: 10212266). These rearrangements regulate many cellular processes,

including cell shape determination, membrane transport, and signal transduction (PubMed:12387735, PubMed:15039356). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:9298994, PubMed:9616160). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse

formation (PubMed:27405666).

Cellular Location Cell membrane; Peripheral membrane protein


{ECO:0000250|UniProtKB:P26041}; Cytoplasmic side


{ECO:0000250 | UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250 | UniProtKB:P26041}. Apical cell membrane {ECO:0000250 | UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 | UniProtKB:P26041}. Cell projection, microvillus membrane {ECO:0000250 | UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 | UniProtKB:P26041}. Cell projection, microvillus {ECO:0000250 | UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250 | UniProtKB:P26041, ECO:0000269 | PubMed:18586956, ECO:0000269 | PubMed:9298994}


Tissue Location


In all tissues and cultured cells studied.

Images









Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.