

# Moesin Rabbit mAb

Catalog # AP76596

## Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	<a href="#">P26038</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	67820

## Additional Information

Gene ID	4478
Other Names	MSN
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A
Format	Liquid

## Protein Information



Name	MSN ( <a href="#">HGNC:7373</a> )
Function	<p>Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:<a href="#">10212266</a>). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (PubMed:<a href="#">10212266</a>). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (PubMed:<a href="#">12387735</a>, PubMed:<a href="#">15039356</a>). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:<a href="#">9298994</a>, PubMed:<a href="#">9616160</a>). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse formation (PubMed:<a href="#">27405666</a>).</p>
Cellular Location	<p>Cell membrane; Peripheral membrane protein {ECO:0000250 UniProtKB:P26041}; Cytoplasmic side {ECO:0000250 UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P26041}. Apical cell membrane {ECO:0000250 UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250 UniProtKB:P26041}; Cytoplasmic side</p>


{ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus membrane  
{ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein  
{ECO:0000250|UniProtKB:P26041}; Cytoplasmic side  
{ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus  
{ECO:0000250|UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250|UniProtKB:P26041, ECO:0000269|PubMed:18586956, ECO:0000269|PubMed:9298994}

**Tissue Location** In all tissues and cultured cells studied.

**Images**

---





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.