10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MTAP Rabbit mAb

Catalog # AP76600

Product Information

Application	WB, IHC-P, FC
Primary Accession	Q13126
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	31236

Additional Information

Gene ID	4507
Other Names	MTAP
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	MTAP {ECO:0000255|HAMAP-Rule:MF_03155}
Synonyms	MSAP
Function	Catalyzes the phosphorolytic cleavage of S-methyl-5'- thioadenosine (MTA) to adenine and 5-methylthioribose1-phosphate (PubMed:3091600, PubMed:8687427, PubMed:10404592). Involved in the breakdown of MTA, a major by-product of polyamine biosynthesis (PubMed:3091600, PubMed:8687427, PubMed:10404592). Responsible for the first step in the methionine salvage pathway after MTA has been generated from S-adenosylmethionine (PubMed:3091600, PubMed:8687427, PubMed:10404592). Has broad substrate specificity with 6-aminopurine nucleosides as preferred substrates (PubMed:10404592).
Cellular Location	Cytoplasm. Nucleus {ECO:0000255|HAMAP- Rule:MF_03155}
Tissue Location	Ubiquitously expressed.

Background

This gene encodes an enzyme that plays a major role in polyamine metabolism and is important for the salvage of both adenine and methionine. The encoded enzyme is deficient in many cancers because this gene and the tumor suppressor p16 gene are co-deleted. Multiple alternatively spliced transcript variants have been described for this gene, but their full-length natures remain unknown. [provided by RefSeq, Jul 2008]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.