

[KD]Olig1 Rabbit mAb

Catalog # AP76633

Product Information

Application	WB, FC
Primary Accession	Q8TAK6
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	27905

Additional Information

Gene ID	116448
Other Names	OLIG1
Dilution	WB~~1:1000 FC~~1:10~50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	OLIG1
Synonyms	BHLHB6, BHLHE21
Function	Orchestrates progenitor cell fates during cortical gliogenesis and gliomagenesis. In cooperation with OLIG2, activates oligodendrocyte precursor cells (OPCs) formation while repressing the generation of olfactory bulb interneuron intermediate progenitors (OBIN) through transcriptional regulation of GSX2. Mechanistically, binds and silences multiple conserved enhancer elements of GSX2. Cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00981}.
Tissue Location	Expressed in the brain, in oligodendrocytes. Strongly expressed in oligodendrogliomas, while expression is weak to moderate in astrocytomas. Expression in glioblastomas is highly variable.

Background

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to be involved in neuron differentiation and regulation of transcription by RNA polymerase II. Predicted to act upstream of or within neuron fate commitment. Predicted to be part of chromatin. Predicted to be active in nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.