

Optineurin Rabbit mAb

Catalog # AP76634

Product Information

ApplicationWB, IHC-PPrimary AccessionQ96CV9ReactivityHumanHostRabbit

Clonality Monoclonal Antibody

Calculated MW 65922

Additional Information

Gene ID 10133

Other Names OPTN

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name OPTN

Function Plays an important role in the maintenance of the Golgi complex, in

membrane trafficking, in exocytosis, through its interaction with myosin VI and Rab8 (PubMed:<u>27534431</u>). Links myosin VI to the Golgi complex and plays an important role in Golgi ribbon formation (PubMed:<u>27534431</u>). Plays a role in the activation of innate immune response during viral infection.

Mechanistically, recruits TBK1 at the Golgi apparatus, promoting its

trans-phosphorylation after RLR or TLR3 stimulation (PubMed:27538435). In turn, activated TBK1 phosphorylates its downstream partner IRF3 to produce IFN-beta/IFNB1. Plays a neuroprotective role in the eye and optic nerve. May act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 and huntingtin (HD). Mediates the interaction of Rab8 with the probable GTPase-activating protein TBC1D17 during

Rab8 with the probable GTPase-activating protein TBC1D17 during Rab8-mediated endocytic trafficking, such as that of transferrin receptor (TFRC/TfR); regulates Rab8 recruitment to tubules emanating from the endocytic recycling compartment (PubMed:22854040). Autophagy receptor that interacts directly with both the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family; targets ubiquitin- coated bacteria

(xenophagy), such as cytoplasmic Salmonella enterica, and appears to

function in the same pathway as SQSTM1 and CALCOCO2/NDP52.


Cellular Location


Cytoplasm, perinuclear region. Golgi apparatus. Golgi apparatus, trans-Golgi network Cytoplasmic vesicle, autophagosome. Cytoplasmic vesicle. Recycling endosome. Note=Found in the perinuclear region and associates with the Golgi apparatus (PubMed:27534431) Colocalizes with MYO6 and RAB8 at the Golgi complex and in vesicular structures close to the plasma membrane. Localizes to LC3-positive cytoplasmic vesicles upon induction of autophagy

Tissue Location

Present in aqueous humor of the eye (at protein level). Expressed in the trabecular meshwork (at protein level) (PubMed:11834836, PubMed:12379221, PubMed:12646749). Expressed in nonpigmented ciliary epithelium (at protein level) (PubMed:11834836) Expressed at high levels in skeletal muscle, also detected in heart, brain, pancreas, kidney, placenta and liver (PubMed:9488477). Expressed in dermal fibroblasts (at protein level) (PubMed:11834836)

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.