

ORAI3 Rabbit mAb

Catalog # AP76635

Product Information

Application	WB, IHC-P
Primary Accession	Q9BRQ5
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	31499

Additional Information

Gene ID	93129
Other Names	ORAI3
Dilution	WB~~1/500-1/1000 IHC-P~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ORAI3
Synonyms	TMEM142C
Function	<p>Pore-forming subunit of two major inward rectifying Ca(2+) channels at the plasma membrane: Ca(2+) release-activated Ca(2+) (CRAC) channels and arachidonate-regulated Ca(2+)-selective (ARC) channels (PubMed:16807233, PubMed:17442569, PubMed:19182790, PubMed:19622606, PubMed:19706554, PubMed:20354224, PubMed:32415068). Assembles with ORAI1 and ORAI2 to form hexameric CRAC channels that mediate Ca(2+) influx upon depletion of endoplasmic reticulum Ca(2+) store and channel activation by Ca(2+) sensor STIM1, a process known as store-operated Ca(2+) entry (SOCE). Various pore subunit combinations may account for distinct CRAC channel spatiotemporal and cell-type specific dynamics. ORAI1 mainly contributes to the generation of Ca(2+) plateaus involved in sustained Ca(2+) entry and is dispensable for cytosolic Ca(2+) oscillations, whereas ORAI2 and ORAI3 generate oscillatory patterns. CRAC channels assemble in Ca(2+) signaling microdomains where Ca(2+) influx is coupled to calmodulin and calcineurin signaling and activation of NFAT transcription factors recruited to ORAI1 via AKAP5. CRAC channels are the main pathway for Ca(2+) influx in T</p>

cells and promote the immune response to pathogens by activating NFAT-dependent cytokine and chemokine transcription (PubMed:[16807233](#), PubMed:[17442569](#), PubMed:[19182790](#), PubMed:[19706554](#), PubMed:[20354224](#), PubMed:[32415068](#)). Assembles with ORAI1 to form channels that mediate store-independent Ca^{2+} influx in response to inflammatory metabolites arachidonate or its derivative leukotriene C₄, termed ARC and LRC channels respectively (PubMed:[19622606](#), PubMed:[32415068](#)).



Cellular Location

Cell membrane; Multi-pass membrane protein. Note=Colocalizes with STIM1 upon store depletion.

Tissue Location

Expressed in both naive and effector T helper cells with higher levels in effector cells.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.