

NCF4 Rabbit mAb

Catalog # AP76641

Product Information


Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	Q15080
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	39032

Additional Information

Gene ID	4689
Other Names	NCF4
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A
Format	Liquid

Protein Information

Name	NCF4 (HGNC:7662)
Synonyms	SH3PXD4
Function	<p>Subunit of the phagocyte NADPH oxidase complex that mediates the transfer of electrons from cytosolic NADPH to O₂ to produce the superoxide anion (O₂(-)) (Probable). In the activated complex, electrons are first transferred from NADPH to flavin adenine dinucleotide (FAD) and subsequently transferred via two heme molecules to molecular oxygen, producing superoxide through an outer-sphere reaction (By similarity). Activation of the NADPH oxidase complex is initiated by the assembly of cytosolic subunits of the NADPH oxidase complex with the core NADPH oxidase complex to form a complex at the plasma membrane or phagosomal membrane (By similarity). This activation process is initiated by phosphorylation dependent binding of the cytosolic NCF1/p47-phox subunit to the C-terminus of CYBA/p22-phox (By similarity).</p>
Cellular Location	<p>Cytoplasm, cytosol. Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Membrane; Peripheral membrane protein.</p> <p>Note=Translocates to the membrane upon activation by phorbol myristate acetate (PMA)</p>
Tissue Location	Expression is restricted to hematopoietic cells.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.