

PBK Rabbit mAb

Catalog # AP76645

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, IHC-P, FC, IP |
| Primary Accession | Q96KB5 |
| Reactivity | Rat, Human, Mouse |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Isotype | IgG |
| Conjugate | Unconjugated |
| Purification | Affinity Purified |
| Calculated MW | 36085 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 55872 |
| Other Names | PBK |
| Dilution | WB~~1:1000 IHC-P~~N/A FC~~1:10~50 IP~~N/A |
| Format | Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

Protein Information

| | |
|------------------------|---|
| Name | PBK |
| Synonyms | TOPK |
| Function | Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage. |
| Tissue Location | Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules. |

Background

This genes encodes a serine/threonine kinase related to the dual specific mitogen-activated protein kinase

kinase (MAPKK) family. Evidence suggests that mitotic phosphorylation is required for its catalytic activity. This mitotic kinase may be involved in the activation of lymphoid cells and support Ticular functions, with a suggested role in the process of spermatogenesis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.