

# PRAS40 Rabbit mAb

Catalog # AP76669

### **Product Information**

ApplicationWB, IPPrimary AccessionQ96B36ReactivityHuman, RatHostRabbit

**Clonality** Monoclonal Antibody

Calculated MW 27383

#### **Additional Information**

**Gene ID** 84335

Other Names AKT1S1

**Dilution** WB~~1/500-1/1000 IP~~N/A

Format Liquid

#### **Protein Information**

Name AKT1S1 {ECO:0000312 | EMBL:AAH16043.1}

**Function** Negative regulator of the mechanistic target of rapamycin complex 1

(mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:17277771, PubMed:17386266, PubMed:17510057, PubMed:29236692). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1 activity by blocking the MTOR substrate- recruitment site (PubMed:29236692). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:17386266, PubMed:18372248). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:16174443,

PubMed: 18372248). May also play a role in nerve growth factor-mediated

neuroprotection (By similarity).

Cellular Location Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the


cytosolic fraction of the brain. {ECO:0000250 | UniProtKB:Q9D1F4}

**Tissue Location** Widely expressed with highest levels of expression in liver and heart.

Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in

normal cell lines (e.g. HEK293)

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.