

Quiescin Q6 Rabbit mAb

Catalog # AP76680

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	O00391
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	82578

Additional Information

Gene ID	5768
Other Names	QSOX1
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50 IP~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	QSOX1
Synonyms	QSCN6 {ECO:0000303 PubMed:9878249}
Function	Catalyzes the oxidation of sulfhydryl groups in peptide and protein thiols to disulfides with the reduction of oxygen to hydrogen peroxide (PubMed: 17331072 , PubMed: 18393449 , PubMed: 23704371 , PubMed: 23867277 , PubMed: 30367560). Plays a role in disulfide bond formation in a variety of extracellular proteins (PubMed: 17331072 , PubMed: 22801504 , PubMed: 23867277 , PubMed: 30367560). In fibroblasts, required for normal incorporation of laminin into the extracellular matrix, and thereby for normal cell-cell adhesion and cell migration (PubMed: 23704371 , PubMed: 23867277 , PubMed: 30367560).
Cellular Location	[Isoform 1]: Golgi apparatus membrane; Single-pass membrane protein. Secreted. Note=A small proportion is secreted, probably via a proteolytic cleavage that removes the membrane anchor

Tissue Location

Expressed in heart, placenta, lung, liver, skeletal muscle, pancreas and very weakly in brain and kidney {ECO:0000269|PubMed:10708601, ECO:0000269|Ref.8}

Background

Catalyzes the oxidation of sulfhydryl groups in peptide and protein thiols to disulfides with the reduction of oxygen to hydrogen peroxide (PubMed:17331072, PubMed:18393449, PubMed:23704371, PubMed:23867277, PubMed:30367560). Plays a role in disulfide bond formation in a variety of extracellular proteins (PubMed:17331072, PubMed:22801504, PubMed:23867277, PubMed:30367560). In fibroblasts, required for normal incorporation of laminin into the extracellular matrix, and thereby for normal cell-cell adhesion and cell migration (PubMed:23704371, PubMed:23867277, PubMed:30367560).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.