

Rad21 Rabbit mAb

Catalog # AP76683

Product Information

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	O60216
Reactivity	Human, Hamster
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	71690

Additional Information

Gene ID	5885
Other Names	RAD21
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

Protein Information




Name	RAD21
Function	[Double-strand-break repair protein rad21 homolog]: As a member of the cohesin complex, involved in sister chromatid cohesion from the time of DNA replication in S phase to their segregation in mitosis, a function that is essential for proper chromosome segregation, post-replicative DNA repair, and the prevention of inappropriate recombination between repetitive regions (PubMed: 11509732). The cohesin complex may also play a role in spindle pole assembly during mitosis (PubMed: 11590136). In interphase, cohesins may function in the control of gene expression by binding to numerous sites within the genome (By similarity). May control RUNX1 gene expression (Probable). Binds to and represses APOB gene promoter (PubMed: 25575569). May play a role in embryonic gut development, possibly through the regulation of enteric neuron development (By similarity).
Cellular Location	[Double-strand-break repair protein rad21 homolog]: Nucleus. Nucleus matrix Chromosome Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=Associates with chromatin (PubMed:11073952, PubMed:11590136). Before prophase, scattered along chromosome arms (PubMed:11073952). During prophase and prometaphase, most cohesins dissociate from the arms of condensing chromosome, possibly through PLK1-mediated phosphorylation (PubMed:11931760). A small amount of cohesin remains in centromeric regions and is removed from chromosomes

only at the onset of anaphase. At anaphase, cleavage by separase/ESPL1 leads to the dissociation of cohesin from chromosomes and chromosome separation (PubMed:11073952, PubMed:11509732)

Tissue Location

Expressed in the gut (at protein level).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.