

SAP97 Rabbit mAb

Catalog # AP76702

Product Information

Application	WB
Primary Accession	Q12959
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	100455

Additional Information

Gene ID	1739
Other Names	DLG1
Dilution	WB~~1:1000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	DLG1 (HGNC:2900)
Function	Essential multidomain scaffolding protein required for normal development (By similarity). Recruits channels, receptors and signaling molecules to discrete plasma membrane domains in polarized cells. Promotes epithelial cell layer barrier function via maintaining cell- cell adhesion (By similarity). May also play a role in adherens junction assembly, signal transduction, cell proliferation, synaptogenesis and lymphocyte activation. Regulates the excitability of cardiac myocytes by modulating the functional expression of Kv4 channels. Functional regulator of Kv1.5 channel. During long-term depression in hippocampal neurons, it recruits ADAM10 to the plasma membrane (PubMed: 23676497).
Cellular Location	Cell membrane; Peripheral membrane protein. Basolateral cell membrane. Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:Q62696}. Postsynaptic density {ECO:0000250 UniProtKB:Q62696}. Synapse {ECO:0000250 UniProtKB:Q62696} Cell membrane, sarcolemma. Apical cell

membrane. Cell junction. Cytoplasm Note=Colocalizes with EPB41 at regions of intercellular contacts Basolateral in epithelial cells (PubMed:12807908). May also associate with endoplasmic reticulum membranes. Mainly found in neurons soma, moderately found at postsynaptic densities (By similarity) {ECO:0000250|UniProtKB:Q62696, ECO:0000269|PubMed:10859302, ECO:0000269|PubMed:12807908, ECO:0000269|PubMed:8922391, ECO:0000269|PubMed:9192623}

Tissue Location

Abundantly expressed in atrial myocardium (at protein level). Expressed in lung fibroblasts, cervical epithelial and B-cells (at protein level). Expressed in the brain (at protein level) (PubMed:23676497). Widely expressed, with isoforms displaying different expression profiles.

Background

Essential multidomain scaffolding protein required for normal development (By similarity). Recruits channels, receptors and signaling molecules to discrete plasma membrane domains in polarized cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.