

SKP2 Rabbit mAb

Catalog # AP76712

Product Information

Application	WB, IHC-P, FC
Primary Accession	Q13309
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	47761

Additional Information

Gene ID	6502
Other Names	SKP2
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	SKP2
Synonyms	FBXL1
Function	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed: 9736735 , PubMed: 11931757 , PubMed: 12435635 , PubMed: 12769844 , PubMed: 12840033 , PubMed: 15342634 , PubMed: 15668399 , PubMed: 15949444 , PubMed: 16103164 , PubMed: 16262255 , PubMed: 16581786 , PubMed: 16951159 , PubMed: 17908926 , PubMed: 17962192 , PubMed: 22464731 , PubMed: 22770219 , PubMed: 32267835). Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition (By similarity). Degradation of CDKN1B/p27kip also requires CKS1 (By similarity). Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, NBN, FOXO1, UBP43, YTHDF2,

and probably MYC, TOB1 and TAL1 (PubMed:[11931757](#), PubMed:[12435635](#), PubMed:[12769844](#), PubMed:[12840033](#), PubMed:[15342634](#), PubMed:[15668399](#), PubMed:[15949444](#), PubMed:[16103164](#), PubMed:[16581786](#), PubMed:[16951159](#), PubMed:[17908926](#), PubMed:[17962192](#), PubMed:[22464731](#), PubMed:[32267835](#)). Degradation of TAL1 also requires STUB1 (PubMed:[17962192](#)). Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2 (PubMed:[9736735](#), PubMed:[16262255](#)). Promotes ubiquitination and destruction of CDH1 in a CK1-dependent manner, thereby regulating cell migration (PubMed:[22770219](#)). Following phosphorylation in response to DNA damage, mediates 'Lys-63'-linked ubiquitination of NBN, promoting ATM recruitment to DNA damage sites and DNA repair via homologous recombination (PubMed:[22464731](#)).

Cellular Location Cytoplasm. Nucleus

Background

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.