

KDM4A Rabbit mAb

Catalog # AP76785

Product Information

ApplicationWB, IPPrimary AccessionO75164HostRabbit

Clonality Monoclonal Antibody

Calculated MW 120662

Additional Information

Gene ID 9682

Other Names KDM4A

Dilution WB~~1/500-1/1000 IP~~N/A

Format Liquid

Protein Information

Name KDM4A

Synonyms | HDM3A, |M|D2, |M|D2A, KIAA0677

Function Histone demethylase that specifically demethylates 'Lys-9' and 'Lys-36'

residues of histone H3, thereby playing a central role in histone code (PubMed:26741168, PubMed:21768309). Does not demethylate histone H3 'Lys-4', H3 'Lys-27' nor H4 'Lys-20'. Demethylates trimethylated H3 'Lys-9' and H3 'Lys-36' residue, while it has no activity on mono- and dimethylated residues. Demethylation of Lys residue generates formaldehyde and succinate. Participates in transcriptional repression of ASCL2 and

E2F-responsive promoters via the recruitment of histone deacetylases and


NCOR1, respectively.

Cellular Location Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00537,

ECO:0000269 | PubMed:15927959, ECO:0000269 | PubMed:16024779}

Tissue Location Ubiquitous..

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.