

KDM4A Rabbit mAb

Catalog # AP76785

Product Information

Application	WB, IP
Primary Accession	O75164
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	120662


Additional Information

Gene ID	9682
Other Names	KDM4A
Dilution	WB~~1/500-1/1000 IP~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

Protein Information

Name	KDM4A
Synonyms	JHDM3A, JMJD2, JMJD2A, KIAA0677
Function	Histone demethylase that specifically demethylates 'Lys-9' and 'Lys-36' residues of histone H3, thereby playing a central role in histone code (PubMed: 26741168 , PubMed: 21768309). Does not demethylate histone H3 'Lys-4', H3 'Lys-27' nor H4 'Lys-20'. Demethylates trimethylated H3 'Lys-9' and H3 'Lys-36' residue, while it has no activity on mono- and dimethylated residues. Demethylation of Lys residue generates formaldehyde and succinate. Participates in transcriptional repression of ASCL2 and E2F-responsive promoters via the recruitment of histone deacetylases and NCOR1, respectively.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00537, ECO:0000269 PubMed:15927959, ECO:0000269 PubMed:16024779}
Tissue Location	Ubiquitous..

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.