

## **CRCP** Rabbit mAb

Catalog # AP76808

## **Product Information**

**Application** WB, IHC-P, IHC-F, IP, ICC

Primary Accession 075575

Reactivity Human, Mouse, Rat

**Host** Rabbit

**Clonality** Monoclonal Antibody

Calculated MW 16871

## **Additional Information**

**Gene ID** 27297

Other Names CRCP

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A

Format Liquid

## **Protein Information**

Name CRCP ( HGNC:17888)

**Function** DNA-dependent RNA polymerase catalyzes the transcription of DNA into

RNA using the four ribonucleoside triphosphates as substrates (PubMed: 20413673, PubMed: 33558764, PubMed: 34675218). Specific peripheric component of RNA polymerase III (Pol III) which synthesizes small

non-coding RNAs including 5S rRNA, snRNAs, tRNAs and miRNAs from at least 500 distinct genomic loci. With POLR3H/RPC8 forms a mobile stalk that protrudes from Pol III core and functions primarily in transcription initiation (By similarity) (PubMed:20413673, PubMed:33558764, PubMed:33558766, PubMed:34675218). Pol III plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type I interferon and NF-kappa-B through the RIG-I pathway (PubMed:19609254,


PubMed: 19631370).


**Cellular Location** Nucleus. Cell membrane {ECO:0000250|UniProtKB:O35427}; Peripheral


membrane protein {ECO:0000250 | UniProtKB:O35427}; Cytoplasmic side


{ECO:0000250 | UniProtKB:O35427}

**Tissue Location** Ubiquitous. Most prevalent in testis.









Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.