10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Neutrophil Elastase Rabbit mAb

Catalog # AP76955

Product Information

Application	WB, IHC-P, IF, FC, ICC
Primary Accession	P08246
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Neutrophil Elastase
Purification	Affinity Chromatography
Calculated MW	28518

Additional Information

Gene ID	1991
Other Names	ELANE
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ELANE
Synonyms	ELA2
Function	Serine protease that modifies the functions of natural killer cells, monocytes and granulocytes (PubMed:15140022). Shows microbicidal activity in neutrophils by mediating cleavage and degradation of bacterial proteins (PubMed:10947984, PubMed:25161283). Capable of killing E.coli but not S.aureus in vitro; digests outer membrane protein A (ompA) in E.coli and K.pneumoniae (PubMed:10947984). Involved for the formation of neutrophil extracellular traps (NETs): translocates into the nucleus downstream of MPO and mediates degadation of histone H4, thereby participating to chromatin decondensation (PubMed:20974816, PubMed:25066128, PubMed:28574339). NETs are mainly composed of DNA fibers and are released by neutrophils to trap pathogens during inflammation (PubMed:20974816, PubMed:25066128, PubMed:28574339). Inhibits C5a-dependent neutrophil enzyme release and

	chemotaxis (PubMed:15140022). Promotes cleavage of GSDMB, thereby inhibiting pyroptosis (PubMed:36899106). Promotes blood coagulation (PubMed:20676107). Through the activation of the platelet fibrinogen receptor integrin alpha-IIb/beta-3, potentiates platelet aggregation induced by a threshold concentration of cathepsin G (CTSG) (PubMed:25211214, PubMed:9111081). Cleaves and thus inactivates tissue factor pathway inhibitor (TFPI) (PubMed:20676107, PubMed:25211214).
Cellular Location	Cytoplasmic vesicle, phagosome. Lysosome Cytolytic granule. Nucleus. Note=Localizes to azurophil granules, a type of primary lysosome found in neutrophils and monocytes that contains antimicrobial proteins (PubMed:20974816, PubMed:25066128) Translocates from azurophil granules to the nucleus downstream of MPO during formation of neutrophil extracellular traps (NETs), and is present in released NETs (PubMed:20974816, PubMed:25066128, PubMed:32434865). Localized in phagolysosomes following ingestion of E.coli by neutrophils (PubMed:10947984).
Tissue Location	Bone marrow cells. Neutrophil (PubMed:10947984).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.