

Niemann Pick C1 Like 1 Rabbit mAb

Catalog # AP76991

Product Information

Application	WB
Primary Accession	Q9UHC9
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human NPC1L1
Purification	Affinity Chromatography
Calculated MW	148728

Additional Information

Gene ID	29881
Other Names	NPC1L1
Dilution	WB~~1/500-1/1000
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	NPC1L1 (HGNC:7898)
Function	Plays a major role in cholesterol homeostasis (PubMed: 22095670). Critical for the uptake of cholesterol across the plasma membrane of the intestinal enterocyte (PubMed: 22095670). Involved in plant sterol absorption, it transports sitosterol, although at lower rates than cholesterol (By similarity). Is the direct molecular target of ezetimibe, a drug that inhibits cholesterol absorption and is approved for the treatment of hypercholesterolemia (PubMed: 15928087). May have a function in the transport of multiple lipids and their homeostasis, thereby influencing lipid metabolism regulation (PubMed: 15671032). May be involved in caveolin trafficking from the plasma membrane (By similarity). In addition, acts as a negative regulator of NPC2 and down-regulates its expression and secretion by inhibiting its maturation and accelerating its degradation (PubMed: 22095670).
Cellular Location	Apical cell membrane; Multi-pass membrane protein. Cell membrane

{ECO:0000250|UniProtKB:Q6T3U3}; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein. Note=Subfractionation of brush border membranes from proximal enterocytes suggests considerable association with the apical membrane fraction. Exists as a predominantly cell surface membrane expressed protein (By similarity). According to PubMed:15671032, localizes in a subcellular vesicular compartment rich in RAB5.

Tissue Location

Widely expressed. Expressed in liver. Also expressed in small intestine, pancreas, kidney, lung, pancreas, spleen, heart, gall bladder, brain, testis, stomach and muscle

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.