

GATAD2A Rabbit mAb

Catalog # AP77007

Product Information

Application	WB, IHC-P, IF, FC, ICC
Primary Accession	Q86YP4
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human p66 alpha
Purification	Affinity Purified
Calculated MW	68063


Additional Information

Gene ID	54815
Other Names	GATAD2A
Dilution	WB~1/500-1/1000 IHC-P~N/A IF~1:50~200 FC~1:10~50 ICC~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	GATAD2A
Function	Transcriptional repressor (PubMed: 12183469 , PubMed: 16415179). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed: 16428440 , PubMed: 28977666). Enhances MBD2-mediated repression (PubMed: 12183469 , PubMed: 16415179). Efficient repression requires the presence of GATAD2B (PubMed: 16415179).
Cellular Location	Nucleus speckle. Nucleus. Chromosome. Note=Speckled nuclear localization requires both CR1 and CR2 regions (PubMed: 16415179). Localizes to sites of DNA damage in a manner partially dependent on ZMYND8 (PubMed: 27732854).
Tissue Location	Ubiquitous, both in fetal and adult tissues.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.