

WRN Rabbit mAb

Catalog # AP77026

Product Information

Application	WB
Primary Accession	Q14191
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human WRN
Purification	Affinity Purified
Calculated MW	162461

Additional Information

Gene ID	7486
Other Names	WRN
Dilution	WB~~1/500-1/1000
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	WRN
Synonyms	RECQ3, RECQL2
Function	Multifunctional enzyme that has magnesium and ATP-dependent 3'-5' DNA-helicase activity on partially duplex substrates (PubMed: 9224595 , PubMed: 9288107 , PubMed: 9611231). Also has 3'→5' exonuclease activity towards double-stranded (ds)DNA with a 5'-overhang (PubMed: 11863428). Has no nuclease activity towards single-stranded (ss)DNA or blunt-ended dsDNA (PubMed: 11863428). Helicase activity is most efficient with (d)ATP, but (d)CTP will substitute with reduced efficiency; strand displacement is enhanced by single-strand binding- protein (heterotrimeric replication protein A complex, RPA1, RPA2, RPA3) (PubMed: 9611231). Binds preferentially to DNA substrates containing alternate secondary structures, such as replication forks and Holliday junctions. May play an important role in the dissociation of joint DNA molecules that can arise as products of homologous recombination, at

stalled replication forks or during DNA repair. Alleviates stalling of DNA polymerases at the site of DNA lesions. Plays a role in the formation of DNA replication focal centers; stably associates with foci elements generating binding sites for RP-A (By similarity). Plays a role in double-strand break repair after gamma- irradiation (PubMed:[9224595](#), PubMed:[9288107](#), PubMed:[9611231](#)). Unwinds some G-quadruplex DNA (d(CGG)n tracts); unwinding seems to occur in both 5'-3' and 3'-5' direction and requires a short single-stranded tail (PubMed:[10212265](#)). d(CGG)n tracts have a propensity to assemble into tetraplex structures; other G-rich substrates from a telomeric or IgG switch sequence are not unwound (PubMed:[10212265](#)). Depletion leads to chromosomal breaks and genome instability (PubMed:[33199508](#)).

Cellular Location

Nucleus, nucleolus. Nucleus. Nucleus, nucleoplasm. Chromosome. Note=Gamma-irradiation leads to its translocation from nucleoli to nucleoplasm and PML regulates the irradiation-induced WRN relocation (PubMed:21639834). Localizes to DNA damage sites (PubMed:27063109).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.