

ICOS Rabbit mAb

Catalog # AP77044

Product Information

Application WB, IF, FC, ICC **Primary Accession** <u>O9Y6W8</u>

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human ICOS

Purification Affinity Purified

Calculated MW 22625

Additional Information

Gene ID 29851

Other Names ICOS

Dilution WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name ICOS

Synonyms AILIM

Function Stimulatory receptor expressed in activated or antigen- experienced T-cells

that plays an important role in the immune response (PubMed: 9930702). Upon binding to its ligand ICOSL expressed on antigen presenting cells (APCs), delivers costimulatory signals that enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines including IL10, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells (PubMed: 33033255). Also acts as a costimulatory receptor critical for the differentiation of T follicular

regulatory cells upon immune challenges such as viral infection

(PubMed:<u>27135603</u>). Mechanistically, potentiates TCR-induced calcium flux by augmenting PLCG1 activation and actin remodeling (By similarity). In addition,

activates PI3K signaling pathways independently of calcium flux

(PubMed: <u>30523347</u>). Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40-mediated class switching of immunoglobin isotypes (By similarity).


Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.