

PARP2 Rabbit mAb

Catalog # AP77112

Product Information

Application	WB, IF, FC, ICC
Primary Accession	Q9UGN5
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human PARP2
Purification	Affinity Chromatography
Calculated MW	66206

Additional Information

Gene ID	10038
Other Names	PARP2
Dilution	WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	PARP2 {ECO:0000303 PubMed:20092359, ECO:0000312 HGNC:HGNC:272}
Function	Poly-ADP-ribosyltransferase that mediates poly-ADP- ribosylation of proteins and plays a key role in DNA repair (PubMed:10364231, PubMed:25043379, PubMed:27471034, PubMed:30104678, PubMed:32028527, PubMed:32939087, PubMed:34108479, PubMed:34486521, PubMed:34874266). Mediates glutamate, aspartate or serine ADP-ribosylation of proteins: the ADP-D-ribosyl group of NAD(+) is transferred to the acceptor carboxyl group of target residues and further ADP-ribosyl groups are transferred to the 2'-position of the terminal adenosine moiety, building up a polymer with an average chain length of 20-30 units (PubMed:25043379, PubMed:30104678, PubMed:30321391). Serine ADP-ribosylation of proteins constitutes the primary form of ADP-ribosylation of proteins in response to DNA damage (PubMed:32939087). Mediates glutamate and aspartate ADP-ribosylation of target proteins in absence of HPF1 (PubMed:25043379). Following interaction with HPF1, catalyzes serine ADP-ribosylation of target

proteins; HPF1 conferring serine specificity by completing the PARP2 active site (PubMed:[28190768](#), PubMed:[32028527](#), PubMed:[34108479](#), PubMed:[34486521](#), PubMed:[34874266](#)). PARP2 initiates the repair of double-strand DNA breaks: recognizes and binds DNA breaks within chromatin and recruits HPF1, licensing serine ADP-ribosylation of target proteins, such as histones, thereby promoting decompaction of chromatin and the recruitment of repair factors leading to the reparation of DNA strand breaks (PubMed:[10364231](#), PubMed:[32939087](#), PubMed:[34108479](#)). HPF1 initiates serine ADP-ribosylation but restricts the polymerase activity of PARP2 in order to limit the length of poly- ADP-ribose chains (PubMed:[34732825](#), PubMed:[34795260](#)). Specifically mediates formation of branched poly-ADP-ribosylation (PubMed:[30104678](#)). Branched poly-ADP-ribose chains are specifically recognized by some factors, such as APLF (PubMed:[30104678](#)). In addition to proteins, also able to ADP-ribosylate DNA: preferentially acts on 5'-terminal phosphates at DNA strand breaks termini in nicked duplex (PubMed:[27471034](#), PubMed:[29361132](#)).


Cellular Location

Nucleus. Chromosome. Note=Recruited to DNA damage sites in a PARP1-dependent process: recognizes and binds poly-ADP-ribose chains produced by PARP1 at DNA damage sites via its N-terminus, leading to its recruitment.

Tissue Location

Widely expressed, mainly in actively dividing tissues (PubMed:[10364231](#)). The highest levels are in the brain, heart, pancreas, skeletal muscle and testis; also detected in kidney, liver, lung, placenta, ovary and spleen; levels are low in leukocytes, colon, small intestine, prostate and thymus (PubMed:[10364231](#))

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.