

Telomeric Repeat Binding Factor 2 Rabbit mAb

Catalog # AP77135

Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	Q15554
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human TRF2
Purification	Affinity Chromatography
Calculated MW	59594

Additional Information

Gene ID	7014
Other Names	TERF2
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	TERF2
Synonyms	TRBF2, TRF2 {ECO:0000303 PubMed:28216226}
Function	Binds the telomeric double-stranded 5'-TTAGGG-3' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes (PubMed: 15608617 , PubMed: 16166375 , PubMed: 20655466 , PubMed: 28216226 , PubMed: 31595153 , PubMed: 9326950 , PubMed: 9326951 , PubMed: 9476899). In addition to its telomeric DNA-binding role, required to recruit a number of factors and enzymes required for telomere protection, including the shelterin complex, TERF2IP/RAP1 and DCLRE1B/Apollo (PubMed: 16166375 , PubMed: 20655466). Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection (PubMed: 16166375). Shelterin associates with arrays of double-stranded 5'-TTAGGG-3' repeats added by telomerase and protects chromosome ends; without its protective activity, telomeres are no longer

hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways (PubMed:[16166375](#)). Together with DCLRE1B/Apollo, plays a key role in telomeric loop (T loop) formation by generating 3' single-stranded overhang at the leading end telomeres: T loops have been proposed to protect chromosome ends from degradation and repair (PubMed:[20655466](#)). Required both to recruit DCLRE1B/Apollo to telomeres and activate the exonuclease activity of DCLRE1B/Apollo (PubMed:[20655466](#), PubMed:[28216226](#)). Preferentially binds to positive supercoiled DNA (PubMed:[15608617](#), PubMed:[20655466](#)). Together with DCLRE1B/Apollo, required to control the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B) needed for telomere replication during fork passage and prevent aberrant telomere topology (PubMed:[20655466](#)). Recruits TERF2IP/RAP1 to telomeres, thereby participating in to repressing homology-directed repair (HDR), which can affect telomere length (By similarity).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00625, ECO:0000269 | PubMed:20655466}. Chromosome, telomere. Note=Colocalizes with telomeric DNA in interphase cells and is located at chromosome ends during metaphase

Tissue Location

Ubiquitous. Highly expressed in spleen, thymus, prostate, uterus, testis, small intestine, colon and peripheral blood leukocytes.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.