10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DHICA Oxidase Rabbit mAb

Catalog # AP77142

Product Information

Application	WB, IHC-P, IP
Primary Accession	P17643
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human TRP1
Purification	Affinity Chromatography
Calculated MW	60724

Additional Information


Gene ID	7306
Other Names	TYRP1
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	TYRP1 (HGNC:12450)
Function	Plays a role in melanin biosynthesis (PubMed: 16704458, PubMed:22556244, PubMed:23504663). Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) (PubMed:28661582). May regulate or influence the type of melanin synthesized (PubMed:16704458, PubMed:22556244). Also to a lower extent, capable of hydroxylating tyrosine and producing melanin (By similarity).
Cellular Location	Melanosome membrane {ECO:0000250|UniProtKB:P07147}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P07147}. Note=Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 {ECO:0000250|UniProtKB:P07147}

Tissue Location	Pigment cells.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.