10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

JAK1 Rabbit mAb

Catalog # AP77164

Product Information

Application	WB
Primary Accession	P23458
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human JAK1
Purification	Affinity Chromatography
Calculated MW	133277

Additional Information

Gene ID	3716
Other Names	JAK1
Dilution	WB~~1/500-1/1000
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	JAK1 (HGNC:6190)
Synonyms	JAK1A, JAK1B
Function	Non-membrane spanning protein tyrosine kinase that phosphorylates signal-transducing subunits of cytokine receptor complexes like IL2RB, IL10RA, IFNAR2, IL6ST, LIFR, OSMR and IL31RA (PubMed:11909529, PubMed:12133952, PubMed:15194700, PubMed:16239216, PubMed:28111307, PubMed:32750333, PubMed:7615558, PubMed:8232552, PubMed:9188471). Functionally, is involved in the IFN-alpha/beta/gamma signal pathway (PubMed:16239216, PubMed:28111307, PubMed:32750333, PubMed:7615558, PubMed:8232552). Mechanistically, in response to interferon-binding to IFNAR1-IFNAR2 heterodimer, phosphorylates and activates its binding partner IFNAR2, creating docking sites for STAT proteins (PubMed:7759950). Directly phosphorylates STAT proteins but also activates STAT signaling through the transactivation of other JAK kinases associated

with signaling receptors (PubMed:16239216, PubMed:32750333, PubMed:8232552). Involved in the MT-RNR2/humanin- mediated signaling pathway leading to STAT3 phosphorylation (PubMed:27384491). Binding of CNTF or the CLCF1/CLF heterodimer to CNTFR leads to IL6ST/gp130-LIFR dimerization followed by activation of JAK1 and JAK2 which in turns phosphorylate IL6ST/gp130 and LIFR (PubMed:11294841). The tyrosine phosphorylated signaling receptors serve in turn as docking proteins for STAT3 (PubMed:11294841). Involved in the oncostatin-M-mediated signaling pathway through both type I OSM receptor complex (heterodimers composed of LIFR and IL6ST) and type II OSM receptor complex (heterodimers composed of OSMR and IL6ST) (PubMed:9188471). Involved in the oncostatin-M-mediated signaling pathway through both type I OSM receptor complex (heterodimers composed of LIFR and IL6ST) and type II OSM receptor complex (heterodimers composed of OSMR and IL6ST) (PubMed:9188471). Involved in the Interleukin-31-mediated signaling pathway through the IL31 receptor complex (heterodimers composed of OSMR and IL31RA) (PubMed:15194700).


Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Wholly intracellular, possibly membrane associated

Tissue Location

Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.