

# SRMS Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7719b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">Q9H3Y6</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB3601
<b>Calculated MW</b>	54507
<b>Antigen Region</b>	456-485

## Additional Information

---

<b>Gene ID</b>	6725
<b>Other Names</b>	Tyrosine-protein kinase Srms, SRMS, C20orf148
<b>Target/Specificity</b>	This SRMS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 456-485 amino acids from the C-terminal region of human SRMS.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SRMS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	SRMS
<b>Synonyms</b>	C20orf148
<b>Function</b>	Non-receptor tyrosine-protein kinase which phosphorylates DOK1 on tyrosine residues (PubMed: <a href="#">23822091</a> ). Also phosphorylates KHDRBS1/SAM68 and VIM on tyrosine residues (PubMed: <a href="#">29496907</a> ). Phosphorylation of

KHDRBS1 is EGF-dependent (PubMed:[29496907](#)). Phosphorylates OTUB1, promoting deubiquitination of RPTOR (PubMed:[35927303](#)).

**Cellular Location**

Cytoplasm. Note=Localizes to punctate cytoplasmic structures.

**Tissue Location**

Highly expressed in most breast cancers (at protein level)


## Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.


## References

Kohmura, N., et al., Mol. Cell. Biol. 14(10):6915-6925 (1994).

## Images




The anti-SRMS Pab (Cat. #AP7719b) is used in Western blot to detect SRMS in mouse kidney tissue lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

SRMS Antibody (P471) (Cat. #AP7719b) western blot analysis in HL-60 cell line lysates (35ug/lane). This



demonstrates the SRMS antibody detected the SRMS protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.