

NR5A2 Rabbit mAb

Catalog # AP77205

Product Information

Application WB
Primary Accession 000482

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human NR5A2

Purification Affinity Chromatography

Calculated MW 61331

Additional Information

Gene ID 2494

Other Names NR5A2

Dilution WB~~1/500-1/1000

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

NR5A2 {ECO:0000303 | PubMed:11595170, ECO:0000312 | HGNC:HGNC:7984}

Function Orphan nuclear receptor that binds DNA as a monomer to the

5'-TCAAGGCCA-3' sequence and controls expression of target genes: regulates key biological processes, such as early embryonic development, cholesterol

and bile acid synthesis pathways, as well as liver and pancreas

morphogenesis (PubMed:16289203, PubMed:18410128, PubMed:21614002, PubMed:32433991, PubMed:38409506, PubMed:9786908). Ligand-binding causes conformational change which causes recruitment of coactivators, promoting target gene activation (PubMed:21614002). The specific ligand is unknown, but specific phospholipids, such as phosphatidylethanolamine, phosphatidylserine, dilauroyl phosphatidylcholine and diundecanoyl phosphatidylcholine can act as ligand in vitro (PubMed:15707893,

PubMed:<u>15723037</u>, PubMed:<u>15897460</u>, PubMed:<u>21614002</u>, PubMed:<u>22504882</u>, PubMed:<u>23737522</u>, PubMed:<u>26416531</u>,

PubMed: <u>26553876</u>). Acts as a pioneer transcription factor, which unwraps

target DNA from histones and elicits local opening of closed chromatin (PubMed:38409506). Plays a central role during preimplantation stages of embryonic development (By similarity). Plays a minor role in zygotic genome activation (ZGA) by regulating a small set of two-cell stage genes (By similarity). Plays a major role in morula development (2-16 cells embryos) by acting as a master regulator at the 8-cell stage, controlling expression of lineage-specifying transcription factors and genes involved in mitosis, telomere maintenance and DNA repair (By similarity). Zygotic NR5A2 binds to both closed and open chromatin with other transcription factors, often at SINE B1/Alu repeats DNA elements, promoting chromatin accessibility at nearby regulatory regions (By similarity). Also involved in the epiblast stage of development and embryonic stem cell pluripotency, by promoting expression of POU5F1/OCT4 (PubMed:27984042). Regulates other processes later in development, such as formation of connective tissue in lower jaw and middle ear, neural stem cell differentiation, ovarian follicle development and Sertoli cell differentiation (By similarity). Involved in exocrine pancreas development and acinar cell differentiation (By similarity). Acts as an essential transcriptional regulator of lipid metabolism (PubMed:20159957). Key regulator of cholesterol 7-alpha- hydroxylase gene (CYP7A) expression in liver (PubMed: 10359768). Also acts as a negative regulator of inflammation in different organs, such as, liver and pancreas (PubMed:20159957). Protects against intestinal inflammation via its ability to regulate glucocorticoid production (By similarity). Plays an anti-inflammatory role during the hepatic acute phase response by acting as a corepressor: inhibits the hepatic acute phase response by preventing dissociation of the N-Cor corepressor complex (PubMed:20159957). Acts as a regulator of immunity by promoting lymphocyte T-cell development, proliferation and effector functions (By similarity). Also involved in resolution of endoplasmic reticulum stress in the liver (By similarity).


Cellular Location

Nucleus. Chromosome

Tissue Location

Abundantly expressed in pancreas, less in liver, very low levels in heart and lung. Expressed in the Hep-G2 cell line (PubMed:9786908). Isoform 1 and isoform 2 seem to be present in fetal and adult liver and Hep-G2 cells (PubMed:10359768)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.